

Test Problem Construction for Single-Objective Bilevel Optimization

Ankur Sinha, Pekka Malo

Department of Information and Service Economy
Aalto University School of Business, Finland
{Firstname.Lastname}@aalto.fi

Kalyanmoy Deb, IEEE Fellow

Department of Mechanical Engineering
Indian Institute of Technology Kanpur, India
deb@iitk.ac.in

KanGAL Report Number 2013002

Abstract

In this paper, we propose a procedure for designing controlled test problems for single-objective bilevel optimization. The proposed procedure is flexible such that the different complexities present in a bilevel problem can be controlled independently of each other. At the same time, the procedure allows to introduce difficulties caused by interaction of the two levels. Using the construction procedure, the paper also provides a test suite of twelve test problems, which consists of eight unconstrained and four constrained problems. The test suite contains problems with scalable variables and constraints, which can be used to evaluate the ability of the algorithms in handling bilevel problems. To provide baseline results, we have solved the proposed test problems using a nested bilevel evolutionary algorithm. The results can be used for comparison, while evaluating the performance of any other bilevel optimization algorithm.

Keywords: Bilevel optimization, bilevel test-suite, test problem construction, evolutionary algorithm.

1 Introduction

Bilevel optimization constitutes a challenging class of optimization problems, where one optimization task is nested within the other. A large number of studies have been conducted in the field of bilevel programming [7, 17, 11, 9], and on its practical applications [2]. Classical approaches commonly used to handle bilevel problems include the Karush-Kuhn-Tucker approach [4, 12], Branch-and-bound techniques [3] and the

use of penalty functions [1]. Despite a significant progress made in classical optimization towards solving bilevel optimization problems, most of these approaches are rendered inapplicable for bilevel problems with higher levels of complexity. Over the last two decades, technological advances and availability of enormous computing resources have given rise to heuristic approaches for solving difficult optimization problems. Heuristics such as evolutionary algorithms are recognized as potent tools for handling challenging classes of optimization problems. A number of studies have been performed towards using evolutionary algorithms [19, 18, 9] for solving bilevel problems. However, the research on evolutionary algorithms for bilevel problems is still in nascent stage, and significant improvement in the existing approaches is required. Most of the heuristic approaches lack a finite time convergence proof for optimization problems. Therefore, it is a common practice among researchers to demonstrate the convergence of their algorithms on a test bed constituting problems with various complexities. To the best knowledge of the authors, in the domain of single objective bilevel programming, there does not exist a systematic framework for constructing bilevel test problems with controlled difficulties. Test problems, which offer various difficulties found in practical application problems, are often required during the construction and evaluation of algorithms.

Past studies [13] on bilevel optimization have introduced a number of simple test problems. However, the levels of difficulty cannot be controlled in these test problems. In most of the studies, the problems are either linear [14], or quadratic [5, 6], or non-scalable with fixed number of decision variables. Application problems in transportation (network design, optimal pricing), economics (Stackelberg games, principal-agent problem, taxation, policy decisions), management (network facility location, coordination of multi-divisional firms), engineering (optimal design, optimal chemical equilibria) etc [10] have also been used to demonstrate the efficiency of algorithms. For most of these problems, the true optimal solution is unknown. Therefore, it is hard to identify, whether a particular solution obtained using an existing approach is close to the optima. Under these uncertainties, it is not possible to systematically evaluate solution procedures on practical problems. These drawbacks pose hurdles in algorithm development, as the performance of the algorithms cannot be evaluated on various difficulty frontiers. A test-suite with controllable level of difficulties helps in understanding the bilevel algorithms better. It gives an information as to what properties of bilevel problems is being efficiently handled by the algorithm and what is not. An algorithm which performs well on the test problem by effectively tackling most of the challenges offered by the test-suite is expected to perform good on other simpler problems as well. Therefore, controlled test problems are necessary for the advancement of the research on bilevel optimization using evolutionary algorithms.

In this paper, we identify the challenges commonly encountered in bilevel optimization problems. A test problem construction procedure is proposed, which mimics these difficulties in a controllable manner. Using the construction procedure, we propose a collection of bilevel test problems scalable in terms of variables and constraints. The proposed scheme allows to control the difficulties at the two levels independently of each other. At the same time, it also allows to control the extent of difficulty arising due to interaction of the two levels. The test problems generated using the framework are such that the optimal solution of the bilevel problem is known. Moreover, the in-

duced set of the bilevel problem is known as a function of the upper level variables. Such information helps the algorithm developers to debug their procedures during the development phase, and also allows to evaluate the convergence properties of the approach.

The paper is organized as follows. In the next section, we explain the structure of a general bilevel optimization problem and introduce the notation that is used throughout the paper. Section 3 presents our framework for constructing scalable test problems for bilevel programming. Thereafter, following the guidelines of the construction procedure, we suggest a set of twelve scalable test problems in Section 4. To create a benchmark for evaluating different solution algorithms, the problems are solved using a simple nested bilevel evolutionary algorithm which is a nested scheme described in Section 5. The results for the baseline algorithm are discussed in Section 6.

2 Description of a Bilevel Problem

A bilevel optimization problem involves two levels of optimization tasks, where one level is nested within the other. The outer optimization task is usually called as upper level optimization task, and the inner optimization task is called as lower level optimization task. The structure of such an optimization problem requires that only the optimal solutions of the inner optimization task are acceptable as feasible members for the outer optimization task. The problem contains two types of variables; namely the upper level variables \mathbf{x}_u , and the lower level variables \mathbf{x}_l . The lower level is optimized with respect to the lower level variables \mathbf{x}_l , and the upper level variables \mathbf{x}_u act as parameters. An optimal lower level vector and the corresponding upper level vector \mathbf{x}_u constitute a feasible upper level solution, provided the upper level constraints are also satisfied. The upper level problem involves all variables $\mathbf{x} = (\mathbf{x}_u, \mathbf{x}_l)$, and the optimization is supposed to be performed with respect to both \mathbf{x}_u and \mathbf{x}_l . In the following, we provide two equivalent formulations for a general bilevel optimization problem with one objective at both levels:

Definition 1 (Bilevel Optimization Problem (BLOP)) *Let $X = X_U \times X_L$ denote the product of the upper-level decision space X_U and the lower-level decision space X_L , i.e. $\mathbf{x} = (\mathbf{x}_u, \mathbf{x}_l) \in X$, if $\mathbf{x}_u \in X_U$ and $\mathbf{x}_l \in X_L$. For upper-level objective function $F : X \rightarrow \mathbb{R}$ and lower-level objective function $f : X \rightarrow \mathbb{R}$, a general bilevel optimization problem is given by*

$$\begin{aligned} & \underset{\mathbf{x} \in X}{\text{Minimize}} && F(\mathbf{x}), \\ & \text{s.t.} && \mathbf{x}_l \in \underset{\mathbf{x}_l \in X_L}{\text{argmin}} \{f(\mathbf{x}) \mid g_i(\mathbf{x}) \geq 0, i \in I\}, \\ & && G_j(\mathbf{x}) \geq 0, j \in J. \end{aligned} \tag{1}$$

where the functions $g_i : X \rightarrow \mathbb{R}$, $i \in I$, represent lower-level constraints and $G_j : X \rightarrow \mathbb{R}$, $j \in J$, is the collection of upper-level constraints.

In the above formulation, a vector $\mathbf{x}^{(0)} = (\mathbf{x}_u^{(0)}, \mathbf{x}_l^{(0)})$ is considered feasible at the upper level, if it satisfies all the upper level constraints, and vector $\mathbf{x}_l^{(0)}$ is optimal

at the lower level for the given $\mathbf{x}_u^{(0)}$. We observe in this formulation that the lower-level problem is a parameterized constraint to the upper-level problem. An equivalent formulation of the bilevel optimization problem is obtained by replacing the lower-level optimization problem with a set value function which maps the given upper-level decision vector to the corresponding set of optimal lower-level solutions. In the domain of Stackelberg games, such mapping is referred as the rational reaction of the follower on the leader's choice \mathbf{x}_u .

Definition 2 (Alternative definition of Bilevel Problem) *Let set-valued function $\Psi : X_U \rightrightarrows X_L$, denote the optimal-solution set mapping of the lower level problem, i.e.*

$$\Psi(\mathbf{x}_u) = \underset{\mathbf{x}_l \in X_L}{\operatorname{argmin}} \{ f(\mathbf{x}) \mid g_i(\mathbf{x}) \geq 0, i \in I \}.$$

A general bilevel optimization problem (BLOP) is then given by

$$\begin{aligned} & \underset{\mathbf{x} \in X}{\operatorname{Minimize}} && F(\mathbf{x}), \\ & \text{s.t.} && \mathbf{x}_l \in \Psi(\mathbf{x}_u), \\ & && G_j(\mathbf{x}) \geq 0, j \in J. \end{aligned} \tag{2}$$

where the function Ψ may be a single-vector valued or a multi-vector valued function depending on whether the lower level function has multiple global optimal solutions or not.

In the test problem construction procedure, the Ψ function provides a convenient description of the relationship between the upper and lower level problems. Figures 1 and 2 illustrate two scenarios, where Ψ can be a single vector valued or a multi-vector valued function respectively. In Figure 1, the lower level problem is shown to be a paraboloid with a single minimum function value corresponding to the set of upper level variables \mathbf{x}_u . Figure 2 represents a scenario where the lower level function is a paraboloid sliced from the bottom with a horizontal plane. This leads to multiple minimum values for the lower level problem, and therefore, multiple lower level solutions correspond to the set of upper level variables \mathbf{x}_u .

3 Test Problem Construction Procedure

The presence of an additional optimization task within the constraints of the upper optimization task leads to a significant increase in complexity, as compared to any single level optimization problem. We identify various kinds of complexities, which a bilevel optimization problem can offer, and provide a test problem construction procedure which can induce these difficulties in a controllable manner. In order to create realistic test problems, the construction procedure should be able to control the scale of difficulties at both levels independently and collectively, such that the performance of algorithms in handling the two levels is evaluated. The test problems created using the construction procedure are expected to be scalable in terms of number of decision variables and constraints, such that the performance of the algorithms can be evaluated




Figure 1: Relationship between upper and lower level variables in case of a single-vector valued mapping. For simplicity the lower level function is in the shape of a paraboloid.




Figure 2: Relationship between upper and lower level variables in case of a multi-vector valued mapping. The lower level function is shown in the shape of a paraboloid with the bottom sliced with a plane.

against increasing number of variables and constraints. The construction procedure should be able to generate test problems with the following properties:

Necessary Properties:

1. The optimal solution of the bilevel optimization should be known.
2. Clear identification of a relationship between the lower level optimal solutions and the upper level variables.

Properties for inducing difficulties:

1. Controlled difficulty in convergence at upper and lower levels.
2. Controlled difficulty caused by interaction of the two levels.
3. Multiple global solutions at the lower level for a given set of upper level variables.
4. Possibility to have either conflict or cooperation between the two levels.
5. Scalability to any number of decision variables at upper and lower levels.
6. Constraints (preferably scalable) at upper and lower levels.

Next, we provide the bilevel test problem construction procedure, which is able to induce most of the difficulties suggested above.

Table 1: Overview of test-problem framework components

Panel A: Decomposition of decision variables			
Upper-level variables		Lower-level variables	
Vector	Purpose	Vector	Purpose
\mathbf{x}_{u1}	Complexity on upper-level	\mathbf{x}_{l1}	Complexity on lower-level
\mathbf{x}_{u2}	Interaction with lower-level	\mathbf{x}_{l2}	Interaction with upper-level

Panel B: Decomposition of objective functions			
Upper-level objective function		Lower-level objective function	
Component	Purpose	Component	Purpose
$F_1(\mathbf{x}_{u1})$	Difficulty in convergence	$f_1(\mathbf{x}_{u1}, \mathbf{x}_{u2})$	Functional dependence
$F_2(\mathbf{x}_{l1})$	Conflict / co-operation	$f_2(\mathbf{x}_{l1})$	Difficulty in convergence
$F_3(\mathbf{x}_{u2}, \mathbf{x}_{l2})$	Difficulty in interaction	$f_3(\mathbf{x}_{u2}, \mathbf{x}_{l2})$	Difficulty in interaction

3.1 Objective functions in the test-problem framework

To create a tractable framework for test-problem construction, we split the upper and lower level functions into three components. Each of the components is specialized for induction of certain kinds of difficulties into the bilevel problem. The functions are determined based on the required complexities at upper and lower levels independently, and also by the required complexities because of the interaction of the two levels. We write a generic bilevel test problem as follows:

$$\begin{aligned}
 F(\mathbf{x}_u, \mathbf{x}_l) &= F_1(\mathbf{x}_{u1}) + F_2(\mathbf{x}_{l1}) + F_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}) \\
 f(\mathbf{x}_u, \mathbf{x}_l) &= f_1(\mathbf{x}_{u1}, \mathbf{x}_{u2}) + f_2(\mathbf{x}_{l1}) + f_3(\mathbf{x}_{u2}, \mathbf{x}_{l2})
 \end{aligned} \tag{3}$$

where

$$\mathbf{x}_u = (\mathbf{x}_{u1}, \mathbf{x}_{u2}) \quad \text{and} \quad \mathbf{x}_l = (\mathbf{x}_{l1}, \mathbf{x}_{l2})$$

In the above equations, each of the levels contains three terms. A summary on the roles of different terms is provided in Table 1. The upper level and lower level variables have been broken into two smaller vectors (see Panel A in Table 1). The vectors \mathbf{x}_{u1} and \mathbf{x}_{l1} are used to induce complexities at the upper and lower levels independently. The vectors \mathbf{x}_{u2} and \mathbf{x}_{l2} are responsible to induce complexities because of interaction. In a similar fashion, we decompose the upper and lower level functions such that each of the components is specialized for a certain purpose only (see Panel B in Table 1). At the upper level, the term $F_1(\mathbf{x}_{u1})$ is responsible for inducing difficulty in convergence solely at the upper level. Similarly, at the lower level, the term $f_2(\mathbf{x}_{l1})$ is responsible for inducing difficulty in convergence solely at the lower level. The term $F_2(\mathbf{x}_{l1})$ decides if there is a conflict or a cooperation between the upper and lower levels. The terms $F_3(\mathbf{x}_{l2}, \mathbf{x}_{u2})$ and $f_3(\mathbf{x}_{l2}, \mathbf{x}_{u2})$ are interaction terms which can be used to induce difficulties because of interaction at the two levels. Term $F_3(\mathbf{x}_{l2}, \mathbf{x}_{u2})$ may also induce a cooperation or a conflict. Finally, $f_1(\mathbf{x}_{u1}, \mathbf{x}_{u2})$ is a fixed term for the lower level optimization problem and does not induce any convergence difficulties. It is used along with the lower level interaction term to create a functional dependence between lower level optimal solution(s) and the upper level variables. The difficulties related to constraints are handled separately.

3.1.1 Controlled difficulty in convergence

The test-problem framework allows to introduce difficulties in terms of convergence at both levels of bilevel optimization problem while retaining sufficient control. To demonstrate this, let us consider the structure of the lower level minimization problem.

For a given $\mathbf{x}_u = (\mathbf{x}_{u1}, \mathbf{x}_{u2})$, the lower level minimization problem is written as

$$\underset{(\mathbf{x}_{l1}, \mathbf{x}_{l2})}{\text{Min}} f(\mathbf{x}_u, \mathbf{x}_l) = f_1(\mathbf{x}_{u1}, \mathbf{x}_{u2}) + f_2(\mathbf{x}_{l1}) + f_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}),$$

where the upper level variables $(\mathbf{x}_{u1}, \mathbf{x}_{u2})$ act as parameters for the optimization problem. The corresponding optimal-set mapping is given by

$$\Psi(\mathbf{x}_u) = \operatorname{argmin}\{f_2(\mathbf{x}_{l1}) + f_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}) : \mathbf{x}_l \in X_L\},$$

where f_1 does not appear due to its independence from \mathbf{x}_l . Since all of the terms are independent of each other, we note that the optimal value of the function f can be recovered by optimizing the functions f_2 and f_3 individually. Function f_2 contains only lower level variables \mathbf{x}_{l1} , which do not interact with upper level variables. It introduces convergence difficulties at the lower level without affecting the upper level optimization task. Function f_3 contains both lower level variables \mathbf{x}_{l2} , and upper level variables \mathbf{x}_{u2} . The optimal value of this function depends on \mathbf{x}_{u2} .

The following example shows that the calibration of the desired difficulty level for the lower level problem boils down to the choice of functions f_2 and f_3 such that their optima are known.

Example 1: To create a simple lower level function, let the dimension of the variable sets be as follows: $\dim(\mathbf{x}_{u1}) = U1$, $\dim(\mathbf{x}_{u2}) = U2$, $\dim(\mathbf{x}_{l1}) = L1$ and $\dim(\mathbf{x}_{l2}) = L2$. Consider a special case where $L2 = U2$, then the three functions could be defined as follows:

$$\begin{aligned} f_1(\mathbf{x}_{u1}, \mathbf{x}_{u2}) &= \sum_{i=1}^{U1} (x_{u1}^i)^2 + \sum_{i=1}^{U2} (x_{u2}^i)^2 \\ f_2(\mathbf{x}_{l1}) &= \sum_{i=1}^{L1} (x_{l1}^i)^2 \\ f_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}) &= \sum_{i=1}^{U2} (x_{u2}^i - x_{l2}^i)^2 \end{aligned}$$

where f_1 affects only the value of the function without inducing any convergence difficulties. The corresponding optimal set mapping Ψ is reduced to an ordinary vector valued function

$$\Psi(\mathbf{x}_u) = \{(\mathbf{x}_{l1}, \mathbf{x}_{l2}) : \mathbf{x}_{l1} = \mathbf{0}, \mathbf{x}_{l2} = \mathbf{x}_{u2}\}.$$

As discussed above, other functions can be chosen with desired complexities to induce difficulties at the lower level and come up with a variety of lower level functions. Similarly, F_1 is a function of \mathbf{x}_{u1} , which does not interact with any lower level variables. It causes convergence difficulties at the upper level without introducing any other form of complexity in the bilevel problem.

3.1.2 Controlled difficulty in interaction

Next, we consider difficulties due to interaction between the upper and lower level optimization tasks. The upper level optimization task is defined as a minimization

problem over the graph of the optimal solution set mapping Ψ , i.e.

$$\text{Min } \{F(\mathbf{x}_u, \mathbf{x}_l) : \mathbf{x}_l \in \Psi(\mathbf{x}_u), \mathbf{x}_u \in X_U\}$$

where the objective function

$$F(\mathbf{x}_u, \mathbf{x}_l) = F_1(\mathbf{x}_{u1}) + F_2(\mathbf{x}_{l1}) + F_3(\mathbf{x}_{u2}, \mathbf{x}_{l2})$$

is a sum of three independent terms. Our primary interest is on the last two terms $F_2(\mathbf{x}_{l1})$ and $F_3(\mathbf{x}_{u2}, \mathbf{x}_{l2})$, which determine the type of interaction there is going to be between the optimization problems. This can be done in two different ways, depending on whether a cooperation or a conflict is desired between the upper and lower level problems.

Definition 3 (Co-operative bilevel test-problem) *A bilevel optimization problem is said to be co-operative, if in the vicinity of \mathbf{x}_l^* for a particular \mathbf{x}_u , an improvement in the lower level function value leads to an improvement in the upper level function value. Within our test problem framework, the independence of terms in the upper level objective function F implies that the co-operative condition is satisfied when for any upper level decision \mathbf{x}_u the corresponding lower level decision $\mathbf{x}_l = (\mathbf{x}_{l1}, \mathbf{x}_{l2})$ is such that $\mathbf{x}_{l1} \in \text{argmin}\{F_2(\mathbf{x}_{l1}) : \mathbf{x}_l \in \Psi(\mathbf{x}_u)\}$ and $\mathbf{x}_{l2} \in \text{argmin}\{F_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}) : \mathbf{x}_l \in \Psi(\mathbf{x}_u)\}$.*

Definition 4 (Conflicting bilevel test-problem) *A bilevel optimization problem is said to be conflicting, if in the vicinity of \mathbf{x}_l^* for a particular \mathbf{x}_u , an improvement in the lower-level function value leads to an adverse effect on the upper level function value. In our framework, a conflicting test problem is obtained when for any upper level decision \mathbf{x}_u the corresponding lower level decision $\mathbf{x}_l = (\mathbf{x}_{l1}, \mathbf{x}_{l2})$ is such that $\mathbf{x}_{l1} \in \text{argmax}\{F_2(\mathbf{x}_{l1}) : \mathbf{x}_l \in \Psi(\mathbf{x}_u)\}$ and $\mathbf{x}_{l2} \in \text{argmax}\{F_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}) : \mathbf{x}_l \in \Psi(\mathbf{x}_u)\}$.*

In the above general form, the functions f_2 and f_3 may have multiple optimal solutions for any given upper level decision \mathbf{x}_u . However, in order to create test problems with tractable interaction patterns, we would like to define them such that each problem has only a single lower level optimum for a given \mathbf{x}_u . To ensure the existence of single lower level optimum, and to enable realistic interactions between the two levels, we consider imposing the following simple restrictions on the objective functions:

Case 1. Creating co-operative interaction: A test problem with co-operative interaction pattern can be created by choosing

$$\begin{aligned} F_2(\mathbf{x}_{l1}) &= f_2(\mathbf{x}_{l1}) \\ F_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}) &= F_4(\mathbf{x}_{u2}) + f_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}), \end{aligned} \tag{4}$$

where $F_4(\mathbf{x}_{u2})$ is any function of \mathbf{x}_{u2} whose minimum is known.

Case 2. Creating conflicting interaction: A test problem with a conflict between the two levels can be created by simply changing the signs of terms f_2 and f_3 on the

right hand side in (4):

$$\begin{aligned} F_2(\mathbf{x}_{l1}) &= -f_2(\mathbf{x}_{l1}) \\ F_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}) &= F_4(\mathbf{x}_{u2}) - f_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}). \end{aligned} \quad (5)$$

The choice of F_2 and F_3 suggested here is a special case, and there can be many other ways to achieve conflict or co-operation using the two functions.

Case 3. Creating mixed interaction: There may be a situation of both cooperation and conflict if functions F_2 and F_3 are chosen with opposite signs as,

$$\begin{aligned} F_2(\mathbf{x}_{l1}) &= f_2(\mathbf{x}_{l1}) \\ F_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}) &= F_4(\mathbf{x}_{u2}) - f_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}) \end{aligned} \quad (6)$$

or

$$\begin{aligned} F_2(\mathbf{x}_{l1}) &= -f_2(\mathbf{x}_{l1}) \\ F_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}) &= F_4(\mathbf{x}_{u2}) + f_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}). \end{aligned} \quad (7)$$

Example 2: Consider a bilevel optimization problem where the lower level task is given by Example 1. According to the above procedures, we can produce a test problem with a conflict between the upper and lower level by defining the upper level objective function as follows:

$$\begin{aligned} F_1(\mathbf{x}_{u1}) &= \sum_{i=1}^{U1} (x_{u1}^i)^2 \\ F_2(\mathbf{x}_{l1}) &= -\sum_{i=1}^{L1} (x_{l1}^i)^2 \\ F_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}) &= -\sum_{i=1}^{U2} (x_{u2}^i - x_{l2}^i)^2. \end{aligned} \quad (8)$$

The chosen formulation corresponds to Case 2, where $F_4(\mathbf{x}_{u2}) = 0$. The final optimal solution of the bilevel problem is $F(\mathbf{x}_u, \mathbf{x}_l) = 0$ for $(\mathbf{x}_u, \mathbf{x}_l) = \mathbf{0}$.

3.1.3 Multiple Global Solutions at Lower Level

In this sub-section, we discuss about constructing test problems with lower level function having multiple global solutions for a given set of upper level variables. To achieve this, we formulate a lower level function which has multiple lower level optima for a given \mathbf{x}_u , such that $\mathbf{x}_l^* \in \Psi(\mathbf{x}_u)$. Then, we ensure that out of all these possible lower level optimal solutions, one of them (\mathbf{x}_l^{**}) corresponds to the best upper level function value, i.e.,

$$\mathbf{x}_l^{**} \in \underset{\mathbf{x}_l^*}{\operatorname{argmin}} \{F(\mathbf{x}_u, \mathbf{x}_l^*) \mid \mathbf{x}_l^* \in \Psi(\mathbf{x}_u)\} \quad (9)$$

To incorporate this difficulty in the problem, we choose the second functions at the upper and lower levels. Given that the term $f_2(\mathbf{x}_{l1})$ is responsible for causing complexities only at the lower level, we can freely formulate it such that it has multiple lower level optimal solutions. From this it necessarily follows that the entire lower level function has multiple optimal solutions.

Example 3: We describe the construction procedure by considering a simple example, where the cardinalities of the variables are, $\dim(\mathbf{x}_{u1}) = 2$, $\dim(\mathbf{x}_{u2}) = 2$, $\dim(\mathbf{x}_{l1}) = 2$ and $\dim(\mathbf{x}_{l2}) = 2$, and the lower level function is defined as follows,

$$\begin{aligned} f_1(\mathbf{x}_{u1}, \mathbf{x}_{u2}) &= (x_{u1}^1)^2 + (x_{u1}^2)^2 + (x_{u2}^1)^2 + (x_{u2}^2)^2 \\ f_2(\mathbf{x}_{l1}) &= (x_{l1}^1 - x_{l1}^2)^2 \\ f_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}) &= (x_{u2}^1 - x_{l2}^1)^2 + (x_{u2}^2 - x_{l2}^2)^2 \end{aligned} \quad (10)$$

Here, we observe that $f_2(\mathbf{x}_{l1})$ induces multiple optimal solutions, as its minimum value is 0 for all $x_{l1}^1 = x_{l1}^2$. At the minimum $f_3(\mathbf{x}_{u2}, \mathbf{x}_{l2})$ fixes the values of x_{l2}^1 and x_{l2}^2 to x_{u2}^1 and x_{u2}^2 respectively. Next, we write the upper level function ensuring that out of the set $x_{l1}^1 = x_{l1}^2$, one of the solutions is best at upper level.

$$\begin{aligned} F_1(\mathbf{x}_{u1}) &= (x_{u1}^1)^2 + (x_{u1}^2)^2 \\ F_2(\mathbf{x}_{l1}) &= (x_{l1}^1)^2 + (x_{l1}^2)^2 \\ F_3(\mathbf{x}_{u2}, \mathbf{x}_{l2}) &= (x_{u2}^1 - x_{l2}^1)^2 + (x_{u2}^2 - x_{l2}^2)^2 \end{aligned} \quad (11)$$

The formulation of $F_2(\mathbf{x}_{l1})$, as sum of squared terms ensures that $x_{l1}^1 = x_{l1}^2 = 0$ provides the best solution at the upper level for any given $(\mathbf{x}_{u1}, \mathbf{x}_{u2})$.

3.2 Difficulties induced by constraints

In this subsection, we discuss about the types of constraints which can be encountered in a bilevel optimization problem. We only consider inequality constraints in this bilevel test problem construction framework. Considering that the bilevel problems have the possibility to have constraints at both levels, and each constraint could be a function of two different kinds of variables, the constrained set at both levels can be further broken down into smaller subsets as follows:

Level	Constraint Set	Subsets	Dependence
Upper	$\mathbf{G} = \{G_j : j \in J\}$	$\mathbf{G} = \mathbf{G}_a \cup \mathbf{G}_b \cup \mathbf{G}_c$	\mathbf{G}_a depends on \mathbf{x}_u \mathbf{G}_b depends on \mathbf{x}_l \mathbf{G}_c depends on \mathbf{x}_u and \mathbf{x}_l
Lower	$\mathbf{g} = \{g_i : i \in I\}$	$\mathbf{g} = \mathbf{g}_a \cup \mathbf{g}_b \cup \mathbf{g}_c$	\mathbf{g}_a depends on \mathbf{x}_u \mathbf{g}_b depends on \mathbf{x}_l \mathbf{g}_c depends on \mathbf{x}_u and \mathbf{x}_l

Table 2: Composition of the constraint sets at both levels.

In Table 2, \mathbf{G} and \mathbf{g} denote the set of constraints at the upper and lower level respectively. Each of the constraint set can be broken into three smaller subsets, as shown in the table. The first subset represents constraints that are functions of the upper level variables only, the second subset represents constraints that are functions of lower level variables only, and the third subset represents constraints that are functions of both upper and lower level variables. The reason for splitting the constraints into smaller subsets is to develop an insight for solving these kinds of problems using an

evolutionary approach. If the first constraint subset (\mathbf{G}_a or \mathbf{g}_a) is non-empty at either of the two levels, then for any given \mathbf{x}_u we should check the feasibility of constraints in the sets \mathbf{G}_a and \mathbf{g}_a , before solving the lower level optimization problem. In case, there is one or more infeasible constraints in \mathbf{g}_a , then the lower level optimization problem does not contain optimal lower level solution (\mathbf{x}_l^*) for the given \mathbf{x}_u . However, if one or more constraints are infeasible within \mathbf{G}_b , then a lower level optimal solution (\mathbf{x}_l^*) may exist for the given \mathbf{x}_u , but the pair $(\mathbf{x}_u, \mathbf{x}_l^*)$ will be infeasible for the bilevel problem. Based on this property, a decision can be made, whether it is useful to solve the lower level optimization problem at all for a given \mathbf{x}_u .

The upper level constraint subsets, \mathbf{G}_b depends on \mathbf{x}_l , and \mathbf{G}_c depends on \mathbf{x}_u and \mathbf{x}_l . The values from these constraints are meaningful only when the lower level vector is an optimal solution to the lower level optimization problem. Based on the various constraints which may be functions of \mathbf{x}_u , or \mathbf{x}_l or both, a bilevel problem introduces different kinds of difficulties in the optimization task. In this paper, we aim to construct such kinds of constrained bilevel test problems which incorporate some of these complexities. We have proposed four constrained bilevel problems, each of which has at least one or more of the following properties,

1. Constraints exist, but are not active at the optimum
2. A subset of constraints, or all the constraints are active at the optimum
3. Upper level constraints are functions of only upper level variables, and lower level constraints are functions of only lower level variables
4. Upper level constraints are functions of upper as well as lower level variables, and lower level constraints are also functions of upper as well as lower level variables
5. Lower level constraints lead to multiple global solutions at the lower level
6. Constraints are scalable at both levels

While describing the test problems in the next section, we discuss the construction procedure for the individual constrained test problems.

4 SMD test problems

By adhering to the design principles introduced in the previous section, we now propose a set of twelve problems which we call as the SMD test problems. Each problem represents a different difficulty level in terms of convergence at the two levels, complexity of interaction between two levels and multi-modalities at each of the levels. The first eight problems are unconstrained and the remaining four are constrained.




Figure 3: Upper and lower level function contours for a four-variable SMD1 test problem.

4.1 SMD1

This is a simple test problem where the lower level problem is a convex optimization task, and the upper level is convex with respect to upper level variables and optimal lower level variables. The two levels cooperate with each other.

$$\begin{aligned}
 F_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\
 F_2 &= \sum_{i=1}^q (x_{l1}^i)^2 \\
 F_3 &= \sum_{i=1}^r (x_{u2}^i)^2 + \sum_{i=1}^r (x_{u2}^i - \tan x_{l2}^i)^2 \\
 f_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\
 f_2 &= \sum_{i=1}^q (x_{l1}^i)^2 \\
 f_3 &= \sum_{i=1}^r (x_{u2}^i - \tan x_{l2}^i)^2
 \end{aligned} \tag{12}$$

The range of variables is as follows,

$$\begin{aligned}
 x_{u1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, p\} \\
 x_{u2}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, r\} \\
 x_{l1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, q\} \\
 x_{l2}^i &\in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right), \quad \forall i \in \{1, 2, \dots, r\}
 \end{aligned} \tag{13}$$

Relationship between upper level variables and lower level optimal variables is given as follows,

$$\begin{aligned}
 x_{l1}^i &= 0, \quad \forall i \in \{1, 2, \dots, p\} \\
 x_{l2}^i &= \tan^{-1} x_{u2}^i, \quad \forall i \in \{1, 2, \dots, r\}
 \end{aligned} \tag{14}$$




Figure 4: Upper level function contours for a four-variable SMD1 test problem.

The values of the variables at the optima are $\mathbf{x}_u = 0$ and \mathbf{x}_l is obtained by the relationship given above. Both the upper and lower level functions are equal to zero at the optima.

Figure 3 shows the contours of the upper and lower level functions with respect to the upper and lower level variables for a four-variable test problem. The problem has two upper level variables and two lower level variables, such that the dimensions of \mathbf{x}_{u1} , \mathbf{x}_{u2} , \mathbf{x}_{l1} and \mathbf{x}_{l2} are all one. Sub-figure P shows the upper level function contours with respect to the upper level variables, assuming that the lower level variables are at the optima. Fixing the upper level variables $(\mathbf{x}_{u1}, \mathbf{x}_{u2})$ at five different locations, i.e. $(2, 2)$, $(-2, 2)$, $(2, -2)$, $(-2, -2)$ and $(0, 0)$, the lower level function contours are shown with respect to the lower level variables. This shows that the contours of the lower level optimization problem may be different for different upper level vectors.

Figure 4 shows the contours of the upper level function with respect to the upper and lower level variables. Sub-figure P once again shows the upper level function contours with respect to the upper level variables. However, sub-figures Q, R, S, T and V now represent the upper level function contours at different $(\mathbf{x}_{u1}, \mathbf{x}_{u2})$, i.e. $(2, 2)$, $(-2, 2)$, $(2, -2)$, $(-2, -2)$ and $(0, 0)$. From sub-figures Q, R, S, T and V, we observe that if the lower level variables move away from its optimal location, the upper level function value deteriorates. This means that the upper level function and the lower level functions are cooperative.




Figure 5: Upper and lower level function contours for a four-variable SMD2 test problem.

4.2 SMD2

This test problem is similar to the SDM1 test problem, however there is a conflict between the upper level and lower level optimization task. The lower level optimization problem is once again a convex optimization task and the upper level optimization is convex with respect to upper level variables and optimal lower level variables. Since, the two levels are conflicting, an inaccurate lower level optimum may lead to upper level function value better than the true optimum for the bilevel problem.

$$\begin{aligned}
 F_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\
 F_2 &= -\sum_{i=1}^q (x_{l1}^i)^2 \\
 F_3 &= \sum_{i=1}^r (x_{u2}^i)^2 - \sum_{i=1}^r (x_{u2}^i - \log x_{l2}^i)^2 \\
 f_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\
 f_2 &= \sum_{i=1}^q (x_{l1}^i)^2 \\
 f_3 &= \sum_{i=1}^r (x_{u2}^i - \log x_{l2}^i)^2
 \end{aligned} \tag{15}$$

The range of variables is as follows,

$$\begin{aligned}
 x_{u1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, p\} \\
 x_{u2}^i &\in [-5, 1], \quad \forall i \in \{1, 2, \dots, r\} \\
 x_{l1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, q\} \\
 x_{l2}^i &\in (0, e], \quad \forall i \in \{1, 2, \dots, r\}
 \end{aligned} \tag{16}$$




Figure 6: Upper level function contours for a four-variable SMD2 test problem.

Relationship between upper level variables and lower level optimal variables is given as follows,

$$\begin{aligned} x_{l1}^i &= 0, \quad \forall i \in \{1, 2, \dots, q\} \\ x_{l2}^i &= \log^{-1} x_{u2}^i, \quad \forall i \in \{1, 2, \dots, r\} \end{aligned} \quad (17)$$

The values of the variables at the optima are $\mathbf{x}_u = 0$ and \mathbf{x}_l is obtained by the relationship given above. Both the upper and lower level functions are equal to zero at the optima.

Figure 5 shows the contours of the upper and lower level functions with respect to the upper and lower level variables for a four-variable test problem. The problem has two upper level variables and two lower level variables, such that the dimension of \mathbf{x}_{u1} , \mathbf{x}_{u2} , \mathbf{x}_{l1} and \mathbf{x}_{l2} are all one. The figure provides the same information about SMD2, as Figure 3 provides about SMD1. However, the shape of the contours differ, which is caused by the use of different F_3 and f_3 functions.

Figure 6 shows the contours of the upper level function with respect to the upper and lower level variables, and provides the same information as Figure 4 provides about SMD1. This figure shows the conflicting nature of the problem caused by using a negative sign in F_2 . The conflicting nature can be observed from the sub-figures Q, R, S, T and U. For a given \mathbf{x}_u , as one moves away from the lower level optimal solution, the upper level function value further reduces. On the other hand, in 5 we observe that moving away from the lower level optimal solution causes an increase in lower level function value.

4.3 SMD3

In this test problem there is a cooperation between the two levels. The difficulty introduced is in terms of multi-modality at the lower level which contains the Rastrigin's function. The upper level is convex with respect to upper level variables and optimal lower level variables.

$$\begin{aligned}
F_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\
F_2 &= \sum_{i=1}^q (x_{l1}^i)^2 \\
F_3 &= \sum_{i=1}^r (x_{u2}^i)^2 + \sum_{i=1}^r ((x_{u2}^i)^2 - \tan x_{l2}^i)^2 \\
f_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\
f_2 &= q + \sum_{i=1}^q \left((x_{l1}^i)^2 - \cos 2\pi x_{l1}^i \right) \\
f_3 &= \sum_{i=1}^r ((x_{u2}^i)^2 - \tan x_{l2}^i)^2
\end{aligned} \tag{18}$$

The range of variables is as follows,

$$\begin{aligned}
x_{u1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, p\} \\
x_{u2}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, r\} \\
x_{l1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, q\} \\
x_{l2}^i &\in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right), \quad \forall i \in \{1, 2, \dots, r\}
\end{aligned} \tag{19}$$

Relationship between upper level variables and lower level optimal variables is given as follows,

$$\begin{aligned}
x_{l1}^i &= 0, \quad \forall i \in \{1, 2, \dots, q\} \\
x_{l2}^i &= \tan^{-1}(x_{u2}^i)^2, \quad \forall i \in \{1, 2, \dots, r\}
\end{aligned} \tag{20}$$

The values of the variables at the optima are $\mathbf{x}_u = 0$ and \mathbf{x}_l is obtained by the relationship given above. Both the upper and lower level functions are equal to zero at the optima. Rastrigin's function used in f_2 has multiple local optima around the global optimum, which introduces convergence difficulties at the lower level.

Sub-figure P in Figure 7 shows the contours of the upper level function with respect to the upper level variables assuming the lower level variables to be optimal at each \mathbf{x}_u . Sub-figures Q, R, S, T, and U show the behavior of the lower level function at 5 different locations of \mathbf{x}_u , which are $(2, 2)$, $(-2, 2)$, $(2, -2)$, $(-2, -2)$ and $(0, 0)$. The problem is once again assumed to have two upper level variables and two lower level variables, such that the dimensions of \mathbf{x}_{u1} , \mathbf{x}_{u2} , \mathbf{x}_{l1} and \mathbf{x}_{l2} are all one. The figure shows that there is a different lower level optimization problem at each \mathbf{x}_u which is required to be solved in order to achieve a feasible solution at the upper level. The contours of the lower level optimization problem differ, based on the location of upper level vector. It can be observed that the rastrigin's function at the lower level introduces multiple local optima into the problem. The contours of the lower level are further distorted because of the presence of the $\tan(\cdot)$ function at the lower level.

In spite of multiple local optima at the lower level, this problem is easier to solve because of the cooperating nature of the functions at the two levels. If a lower level optimization problem is stuck at a local optimum for a particular \mathbf{x}_u (say $\mathbf{x}_u^{(0)}$), then it will have a poorer objective function value at the upper level. However, as soon as another lower level optimization problem is solved in the vicinity of $\mathbf{x}_u^{(0)}$, which attains




Figure 7: Upper and lower level function contours for a four-variable SMD3 test problem.

a global optimum, then it will have a better objective function value at the upper level, and will dominate the previous inaccurate solution. Therefore, a method which is able to handle multi-modality at the lower level at least in few of its lower level optimization runs will be able to successfully solve this problem.

4.4 SMD4

In this test problem there is a conflict between the two levels. The difficulty is in terms of multi-modality at the lower level which once again contains the Rastrigin's function. The upper level is convex with respect to upper level variables and optimal lower level variables.

$$\begin{aligned}
 F_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\
 F_2 &= -\sum_{i=1}^q (x_{l1}^i)^2 \\
 F_3 &= \sum_{i=1}^r (x_{u2}^i)^2 - \sum_{i=1}^r (|x_{u2}^i| - \log(1 + x_{l2}^i))^2 \\
 f_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\
 f_2 &= q + \sum_{i=1}^q ((x_{l1}^i)^2 - \cos 2\pi x_{l1}^i) \\
 f_3 &= \sum_{i=1}^r (|x_{u2}^i| - \log(1 + x_{l2}^i))^2
 \end{aligned} \tag{21}$$

The range of variables is as follows,

$$\begin{aligned}
 x_{u1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, p\} \\
 x_{u2}^i &\in [-1, 1], \quad \forall i \in \{1, 2, \dots, r\} \\
 x_{l1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, q\} \\
 x_{l2}^i &\in [0, e], \quad \forall i \in \{1, 2, \dots, r\}
 \end{aligned} \tag{22}$$




Figure 8: Upper and lower level function contours for a four-variable SMD4 test problem.

Relationship between upper level variables and lower level optimal variables is given as follows,

$$\begin{aligned} x_{l1}^i &= 0, \quad \forall i \in \{1, 2, \dots, q\} \\ x_{l2}^i &= \log^{-1} |x_{u2}^i| - 1, \quad \forall i \in \{1, 2, \dots, r\} \end{aligned} \quad (23)$$

The values of the variables at the optima are $\mathbf{x}_u = 0$ and \mathbf{x}_l is obtained by the relationship given above. Both the upper and lower level functions are equal to zero at the optima.

Figure 8 represents the same information as in Figure 7 for a four-variable bilevel problem. However, this problem involves conflict between the two levels, which make the problem significantly more difficult than the previous test problem. For this test problem if a lower level optimization problem is stuck at a local optimum for a particular \mathbf{x}_u , then it will end up having a better objective function value at the upper level, than what it will attain at the true global lower level optimum. Therefore, even if another lower level optimization problem is successfully solved in the vicinity of \mathbf{x}_u , the previous inaccurate solution will dominate the new solution at the upper level. This problem can be handled only by those methods which are able to efficiently handle lower level multi-modality without getting stuck in a local basin.

4.5 SMD5

In this test problem, there is a conflict between the two levels. The difficulty introduced is in terms of multi-modality and convergence at the lower level. The lower level problem contains the Rosenbrock's (banana) function such that the global optimum

lies in a long, narrow, flat parabolic valley. The upper level is convex with respect to upper level variables and optimal lower level variables.

$$\begin{aligned}
F_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\
F_2 &= -\sum_{i=1}^q \left((x_{l1}^{i+1} - (x_{l1}^i)^2) + (x_{l1}^i - 1)^2 \right) \\
F_3 &= \sum_{i=1}^r (x_{u2}^i)^2 - \sum_{i=1}^r (|x_{u2}^i| - (x_{l2}^i)^2)^2 \\
f_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\
f_2 &= \sum_{i=1}^q \left((x_{l1}^{i+1} - (x_{l1}^i)^2) + (x_{l1}^i - 1)^2 \right) \\
f_3 &= \sum_{i=1}^r (|x_{u2}^i| - (x_{l2}^i)^2)^2
\end{aligned} \tag{24}$$

The range of variables is as follows,

$$\begin{aligned}
x_{u1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, p\} \\
x_{u2}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, r\} \\
x_{l1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, q\} \\
x_{l2}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, r\}
\end{aligned} \tag{25}$$

Relationship between upper level variables and lower level optimal variables is given as follows,

$$\begin{aligned}
x_{l1}^i &= 1, \quad \forall i \in \{1, 2, \dots, q\} \\
x_{l2}^i &= \sqrt{|x_{u2}^i|}, \quad \forall i \in \{1, 2, \dots, r\}
\end{aligned} \tag{26}$$

The values of the variables at the optima are $\mathbf{x}_u = 0$ and \mathbf{x}_l is obtained by the relationship given above. Both the upper and lower level functions are equal to zero at the optima.

4.6 SMD6

In this test problem, once again there is a conflict between the two levels. This problem is different from previous problems such that it contains infinitely many global solutions at the lower level, for any given upper level vector. Out of the entire global solution set, there is only a single lower level point which corresponds to the best upper level function value.

$$\begin{aligned}
F_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\
F_2 &= -\sum_{i=1}^q (x_{l1}^i)^2 + \sum_{i=q+1}^{q+s} (x_{l1}^i)^2 \\
F_3 &= \sum_{i=1}^r (x_{u2}^i)^2 - \sum_{i=1}^r (x_{u2}^i - x_{l2}^i)^2 \\
f_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\
f_2 &= \sum_{i=1}^q (x_{l1}^i)^2 + \sum_{i=q+1, i=i+2}^{q+s-1} (x_{l1}^{i+1} - x_{l1}^i)^2 \\
f_3 &= \sum_{i=1}^r (x_{u2}^i - x_{l2}^i)^2
\end{aligned} \tag{27}$$

The range of variables is as follows,

$$\begin{aligned}
x_{u1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, p\} \\
x_{u2}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, r\} \\
x_{l1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, q+s\} \\
x_{l2}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, r\}
\end{aligned} \tag{28}$$




Figure 9: Plot of the term in f_2 responsible for creating multiple optimum solutions at the lower level. The value of the term is zero at all the points in the valley.

Relationship between upper level variables and lower level optimal variables is given as follows,

$$\begin{aligned} x_{l1}^i &= 0, \quad \forall i \in \{1, 2, \dots, q\} \\ x_{l2}^i &= x_{u2}^i, \quad \forall i \in \{1, 2, \dots, r\} \end{aligned} \quad (29)$$

The values of the variables at the optima are $\mathbf{x}_u = 0$ and \mathbf{x}_l is obtained by the relationship given above. Both the upper and lower level functions are equal to zero at the optima.

Figure 9 shows the second term $((x_{l1}^i - x_{l1}^j)^2)$, for $s = 2$) for function f_2 , and its contours at the lower level. It could be observed from the figure that all the points along $x_{l1}^j = x_{l1}^i$ have a value 0 for the function f_2 . All these points are responsible for introducing multiple global optimal solutions at the lower level for any given upper level variable vector. However, out of all the global optimal solutions at the lower level, the solution $x_{l1}^j = x_{l1}^i = 0$ provides the best function value at the upper level for any given upper level variable vector.

4.7 SMD7

In this test problem, we introduce complexities at the upper level, while keeping the lower level optimization task relatively simpler. Most of the previous test problems would be useful to test the ability of algorithms in handling lower level optimization task efficiently. However, this test problem contains multi-modality at the upper level, which demands a global optimization approach at the upper level. The function F_1 at the upper level represents a slightly modified Griewank function.

$$\begin{aligned}
F_1 &= 1 + \frac{1}{400} \sum_{i=1}^p (x_{u1}^i)^2 - \prod_{i=1}^p \left(\cos \frac{x_{u1}^i}{\sqrt{i}} \right) \\
F_2 &= - \sum_{i=1}^q (x_{l1}^i)^2 \\
F_3 &= \sum_{i=1}^r (x_{u2}^i)^2 - \sum_{i=1}^r (x_{u2}^i - \log x_{l2}^i)^2 \\
f_1 &= \sum_{i=1}^p (x_{u1}^i)^3 \\
f_2 &= \sum_{i=1}^q (x_{l1}^i)^2 \\
f_3 &= \sum_{i=1}^r (x_{u2}^i - \log x_{l2}^i)^2
\end{aligned} \tag{30}$$

The range of variables is as follows,

$$\begin{aligned}
x_{u1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, p\} \\
x_{u2}^i &\in [-5, 1], \quad \forall i \in \{1, 2, \dots, r\} \\
x_{l1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, q\} \\
x_{l2}^i &\in (0, e], \quad \forall i \in \{1, 2, \dots, r\}
\end{aligned} \tag{31}$$

Relationship between upper level variables and lower level optimal variables is given as follows,

$$\begin{aligned}
x_{l1}^i &= 0, \quad \forall i \in \{1, 2, \dots, q\} \\
x_{l2}^i &= \log^{-1} x_{u2}^i, \quad \forall i \in \{1, 2, \dots, r\}
\end{aligned} \tag{32}$$

The values of the variables at the optima are $\mathbf{x}_u = 0$ and \mathbf{x}_l is obtained by the relationship given above. Both the upper and lower level functions are equal to zero at the optima.

4.8 SMD8

This test problem once again tests the ability of the algorithms in handling multi-modality at the upper level, and handling a complex lower level problem at the same time. There is also a conflict between the upper level and lower level optimization tasks. The lower level objective contains the Rosenbrock's (banana) function, and the upper level objective contains the multi-modal Ackley's function.

$$\begin{aligned}
F_1 &= 20 + e - 20 \exp \left(-0.2 \sqrt{\frac{1}{p} \sum_{i=1}^p (x_{u1}^i)^2} \right) - \exp \left(\frac{1}{p} \sum_{i=1}^p \cos 2\pi x_{u1}^i \right) \\
F_2 &= - \sum_{i=1}^q \left((x_{l1}^{i+1} - (x_{l1}^i)^2) + (x_{l1}^i - 1)^2 \right) \\
F_3 &= \sum_{i=1}^r (x_{u2}^i)^2 - \sum_{i=1}^r (x_{u2}^i - (x_{l2}^i)^3)^2 \\
f_1 &= \sum_{i=1}^p |x_{u1}^i| \\
f_2 &= \sum_{i=1}^q \left((x_{l1}^{i+1} - (x_{l1}^i)^2) + (x_{l1}^i - 1)^2 \right) \\
f_3 &= \sum_{i=1}^r (x_{u2}^i - (x_{l2}^i)^3)^2
\end{aligned} \tag{33}$$

The range of variables is as follows,

$$\begin{aligned}
x_{u1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, p\} \\
x_{u2}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, r\} \\
x_{l1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, q\} \\
x_{l2}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, r\}
\end{aligned} \tag{34}$$

Relationship between upper level variables and lower level optimal variables is given as follows,

$$\begin{aligned} x_{l1}^i &= 0, \quad \forall i \in \{1, 2, \dots, q\} \\ x_{l2}^i &= (x_{u2}^i)^{\frac{1}{3}}, \quad \forall i \in \{1, 2, \dots, r\} \end{aligned} \quad (35)$$

The values of the variables at the optima are $\mathbf{x}_u = 0$ and \mathbf{x}_l is obtained by the relationship given above. Both the upper and lower level functions are equal to zero at the optima.

4.9 SMD9

In this test problem, we introduce constraints at both upper and lower levels. Constraints are defined such that they causes convergence difficulties at both levels independently. One constraint is introduced at each level, such that the upper level constraint is a function of the upper level variables, and the lower level constraint is a function of the lower level variables. The constraints divide the search space into annular regions, and causes convergence difficulties without altering the global optimum. The constraint at the upper as well as the lower level is however, inactive at the optimum. The two levels are once again conflicting in nature, such that, an inaccurate lower level optimum may lead to upper level function value better than the true optimum for the bilevel problem.

$$\begin{aligned} F_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\ F_2 &= -\sum_{i=1}^q (x_{l1}^i)^2 \\ F_3 &= \sum_{i=1}^r (x_{u2}^i)^2 - \sum_{i=1}^r (x_{u2}^i - \log(1 + x_{l2}^i))^2 \\ f_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\ f_2 &= \sum_{i=1}^q (x_{l1}^i)^2 \\ f_3 &= \sum_{i=1}^r (x_{u2}^i - \log(1 + x_{l2}^i))^2 \end{aligned} \quad (36)$$

The upper and lower level constraints are as follows:

Upper level constraint

$$G_1 : \frac{\sum_{i=1}^p (x_{u1}^i)^2 + \sum_{i=1}^r (x_{u2}^i)^2}{a} - \left[\frac{\sum_{i=1}^p (x_{u1}^i)^2 + \sum_{i=1}^r (x_{u2}^i)^2}{a} + \frac{0.5}{b} \right] \geq 0$$

Lower level constraint

$$g_1 : \frac{\sum_{i=1}^p (x_{l1}^i)^2 + \sum_{i=1}^r (x_{l2}^i)^2}{a} - \left[\frac{\sum_{i=1}^p (x_{l1}^i)^2 + \sum_{i=1}^r (x_{l2}^i)^2}{a} + \frac{0.5}{b} \right] \geq 0 \quad (37)$$

Where $a = 1$ and $b = 1$.

The range of variables is as follows,

$$\begin{aligned} x_{u1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, p\} \\ x_{u2}^i &\in [-5, 1], \quad \forall i \in \{1, 2, \dots, r\} \\ x_{l1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, q\} \\ x_{l2}^i &\in (-1, -1 + \epsilon], \quad \forall i \in \{1, 2, \dots, r\} \end{aligned} \quad (38)$$

Relationship between upper level variables (feasible with respect to upper level constraints) and lower level optimal variables is given as follows,

$$\begin{aligned} x_{l1}^i &= 0, \quad \forall i \in \{1, 2, \dots, q\} \\ x_{l2}^i &= \log^{-1} x_{u2}^i - 1, \quad \forall i \in \{1, 2, \dots, r\} \end{aligned} \quad (39)$$




Figure 10: Feasible and infeasible regions in case of a two-variable constraint function.

Figure 10 shows the restricted search space for the upper level optimization task when it is a function of two upper level variables, i.e. $p = 1$ and $r = 1$. The search space looks similar at the lower level when $q = 1$ and $r = 1$. For higher number of variables, the annular rings transform into spherical shells. The values of the variables at the optima are $x_u = 0$ and $x_l = 0$. Both the upper and lower level functions are equal to zero at the optima.

4.10 SMD10

In this test problem, we introduce constraints at the upper as well as the lower level such that they are scalable. As the number of variables are varied at the upper and the lower levels, the number of constraints also vary. This is different from the previous problem such that all the constraints are active at the optimum. However, in this function once again we have the upper level constraints as functions of the upper level variables, and the lower level constraints as functions of the lower level variables.

$$\begin{aligned}
 F_1 &= \sum_{i=1}^p (x_{u1}^i - 2)^2 \\
 F_2 &= \sum_{i=1}^q (x_{l1}^i)^2 \\
 F_3 &= \sum_{i=1}^r (x_{u2}^i - 2)^2 - \sum_{i=1}^r (x_{u2}^i - \tan x_{l2}^i)^2 \\
 f_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\
 f_2 &= \sum_{i=1}^q (x_{l1}^i - 2)^2 \\
 f_3 &= \sum_{i=1}^r (x_{u2}^i - \tan x_{l2}^i)^2
 \end{aligned} \tag{40}$$

The upper and lower level constraints are as follows:

Upper level constraints

$$\begin{aligned} G_j &: x_{u1}^j - \sum_{i=1, i \neq j}^p (x_{u1}^i)^3 - \sum_{i=1}^r (x_{u2}^i)^3 \geq 0, \quad \forall j \in \{1, 2, \dots, p\} \\ G_{p+j} &: x_{u2}^j - \sum_{i=1, i \neq j}^r (x_{u2}^i)^3 - \sum_{i=1}^p (x_{u1}^i)^3 \geq 0, \quad \forall j \in \{1, 2, \dots, r\} \end{aligned} \quad (41)$$

Lower level constraints

$$g_j : x_{l1}^j - \sum_{i=1, i \neq j}^q (x_{l1}^i)^3 \geq 0, \quad \forall j \in \{1, 2, \dots, q\}$$

The range of variables is as follows,

$$\begin{aligned} x_{u1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, p\} \\ x_{u2}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, r\} \\ x_{l1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, q\} \\ x_{l2}^i &\in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right), \quad \forall i \in \{1, 2, \dots, r\} \end{aligned} \quad (42)$$

Relationship between upper level variables (feasible with respect to upper level constraints) and lower level optimal variables is given as follows,

$$\begin{aligned} x_{l1}^i &= \frac{1}{\sqrt{q-1}}, \quad \forall i \in \{1, 2, \dots, q\} \\ x_{l2}^i &= \tan^{-1} x_{u2}^i, \quad \forall i \in \{1, 2, \dots, r\} \end{aligned} \quad (43)$$

The values of the variables at the optima are $\mathbf{x}_u = \frac{1}{\sqrt{p+r-1}}$, and \mathbf{x}_l is obtained by the relationship given above.

Figure 11 shows the feasible region of the search space for the upper level optimization task, when the upper level objective it is a function of two upper variables, i.e. $p = 1, r = 1$. The shaded part in the figure shows the feasible region, and the dotted lines show the contours of the upper level objective function. For such a two variable upper level objective function, the optima lies at one of the intersections $((\mathbf{x}_{u1}, \mathbf{x}_{u2}) = (1, 1))$ of the constraints shown in the figure.

4.11 SMD11

In this test problem, we introduce constraints which are functions of upper as well as lower variables at both levels. The constraints at the upper level are scalable, but there is just a single constraint at the lower level. The constraint at the lower level introduces multiple global optimal solutions at the lower level for any given upper level vector. At the optimum of the bilevel problem, the lower level constraint is active as well as the upper level constraints are active. The upper level constraints eliminate a large part of the global optimal solutions from the lower level.

$$\begin{aligned} F_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\ F_2 &= -\sum_{i=1}^q (x_{l1}^i)^2 \\ F_3 &= \sum_{i=1}^r (x_{u2}^i)^2 - \sum_{i=1}^r (x_{u2}^i - \log x_{l2}^i)^2 \\ f_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\ f_2 &= \sum_{i=1}^q (x_{l1}^i)^2 \\ f_3 &= \sum_{i=1}^r (x_{u2}^i - \log x_{l2}^i)^2 \end{aligned} \quad (44)$$




Figure 11: Feasible and infeasible regions in case of a two-variable constraint function.

The upper and lower level constraints are as follows:

$$\begin{aligned}
 &\text{Upper level constraints} \\
 &G_j : x_{u2}^j \geq \frac{1}{\sqrt{r}} + \log x_{l2}^j, \quad \forall j \in \{1, 2, \dots, r\} \\
 &\text{Lower level constraint} \\
 &g_1 : \sum_{i=1}^r (x_{u2}^i - \log x_{l2}^i)^2 \geq 1
 \end{aligned} \tag{45}$$

The range of variables is as follows,

$$\begin{aligned}
 x_{u1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, p\} \\
 x_{u2}^i &\in [-1, 1], \quad \forall i \in \{1, 2, \dots, r\} \\
 x_{l1}^i &\in [-5, 10], \quad \forall i \in \{1, 2, \dots, q\} \\
 x_{l2}^i &\in \left[\frac{1}{e}, e\right], \quad \forall i \in \{1, 2, \dots, r\}
 \end{aligned} \tag{46}$$

Relationship between upper level variables and lower level optimal variables is given as follows,

$$\begin{aligned}
 x_{l1}^i &= 0, \quad \forall i \in \{1, 2, \dots, q\} \\
 \mathbf{x}_{l2} &: \sum_{i=1}^r (x_{u2}^i - \log x_{l2}^i)^2 = 1
 \end{aligned} \tag{47}$$

The values of the variables at the optima are $\mathbf{x}_{u1} = 0$, $\mathbf{x}_{u2} = 0$, $\mathbf{x}_{l1} = 0$, and $\mathbf{x}_{l2} = \log^{-1} \frac{-1}{\sqrt{r}}$. The upper level function value is -1 and the lower level function value is $+1$ at the optima.

Figure 12 shows the constraints at the upper as well as the lower level when $r = 2$. In this example, there is one constraint at the lower level and two constraints at the upper level. All the solutions on the lower level constraint represent optimal solutions to the lower level f_3 . When $\mathbf{x}_{l1} = 0$, such that the function f_2 is also optimal, the solutions on the constraint are optimal solutions to the lower level problem for a given \mathbf{x}_u . It can be observed that the two constraints at the upper level eliminate all the lower level optimal solutions except one. The figure shows feasible region with respect to




Figure 12: Feasible and infeasible regions of SMD11 for a particular upper level vector.

upper level constraints for the upper level problem. However, only point \mathbf{p} represents a feasible solution for the upper level problem for a given \mathbf{x}_u , as it is the lower level optimal solution lying in the upper level constraint feasible region. This problem differs from SDM6, which also contained multiple global solutions at the lower level, in two ways. Firstly, multiple global solutions at the lower level are introduced by lower level constraints in this problem, whereas in the previous problem, it was the lower level objective function which was entirely responsible for introducing multiple global solutions. Secondly, out of the multiple global solutions from the lower level, a single solution is selected based on upper level constraints, whereas in the previous problem all the lower level global solutions were feasible, but one of those solutions had the best upper level objective value.

4.12 SMD12

This test problem is a combination of the previous two test problems, and involves a number of difficulties. The test problem has scalable constraints at both levels, and the constraints are a function of both upper as well as lower level variables. At the same time, any lower level optimization problem for a given set of upper level variables has multiple global optima. All the lower level constraints are active at the bilevel optimum.

$$\begin{aligned}
 F_1 &= \sum_{i=1}^p (x_{u1}^i - 2)^2 \\
 F_2 &= \sum_{i=1}^q (x_{l1}^i)^2 \\
 F_3 &= \sum_{i=1}^r (x_{u2}^i - 2)^2 + \sum_{i=1}^r \tan |x_{l2}^i| - \sum_{i=1}^r (x_{u2}^i - \tan x_{l2}^i)^2 \\
 f_1 &= \sum_{i=1}^p (x_{u1}^i)^2 \\
 f_2 &= \sum_{i=1}^q (x_{l1}^i - 2)^2 \\
 f_3 &= \sum_{i=1}^r (x_{u2}^i - \tan x_{l2}^i)^2
 \end{aligned} \tag{48}$$

The upper and lower level constraints are as follows:

$$\begin{aligned}
& \text{Upper level constraints} \\
& x_{u2}^i - \tan x_{l2}^i \geq 0, \quad \forall i \in \{1, 2, \dots, r\} \\
& x_{u1}^j - \sum_{i=1, i \neq j}^p (x_{u1}^i)^3 - \sum_{i=1}^r (x_{u2}^i)^3 \geq 0, \quad \forall j \in \{1, 2, \dots, p\} \\
& x_{u2}^j - \sum_{i=1, i \neq j}^r (x_{u2}^i)^3 - \sum_{i=1}^p (x_{u1}^i)^3 \geq 0, \quad \forall j \in \{1, 2, \dots, r\} \\
& \text{Lower level constraints} \\
& \sum_{i=1}^r (x_{u2}^i - \tan x_{l2}^i)^2 \geq 1 \\
& x_{l1}^j - \sum_{i=1, i \neq j}^p (x_{l1}^i)^3, \quad \forall j \in \{1, 2, \dots, q\}
\end{aligned} \tag{49}$$

The range of variables is as follows,

$$\begin{aligned}
x_{u1}^i & \in [-5, 10], \quad \forall i \in \{1, 2, \dots, p\} \\
x_{u2}^i & \in [-14.10, 14.10], \quad \forall i \in \{1, 2, \dots, r\} \\
x_{l1}^i & \in [-5, 10], \quad \forall i \in \{1, 2, \dots, q\} \\
x_{l2}^i & \in (-1.5, 1.5), \quad \forall i \in \{1, 2, \dots, r\}
\end{aligned} \tag{50}$$

Relationship between upper level variables and lower level optimal variables is given as follows,

$$\begin{aligned}
x_{l1}^i & = \frac{1}{\sqrt{q-1}}, \quad \forall i \in \{1, 2, \dots, q\} \\
\mathbf{x}_{l2} & : \sum_{i=1}^r (x_{u2}^i - \tan x_{l2}^i)^2 = 1
\end{aligned} \tag{51}$$

The values of the variables at the optima are $\mathbf{x}_{u1} = \frac{1}{\sqrt{p+r-1}}$, $\mathbf{x}_{u2} = \frac{1}{\sqrt{p+r-1}}$, $\mathbf{x}_{l1} = \sqrt{q-1}$, and $\mathbf{x}_{l2} = \tan^{-1}(\frac{1}{\sqrt{p+r-1}} - \frac{1}{\sqrt{r}})$.

4.13 Summary

The properties of the SMD test problems are summarized in Table 3. In the table, $N = \text{No}$ and $Y = \text{Yes}$. It can be observed that the 12 test problems are a good mix of various difficulties, which we discussed in the prior sections. We have tried to put the problems in an increasing order of difficulty. The last test problem can be observed to contain most of the difficulties except multi-modalities. This table will be helpful in testing algorithms for bilevel optimization. For example, if a new algorithm is able to solve SMD1 but not SMD2, one readily concludes that the algorithm is unable to handle a conflict. Similarly, if the algorithm is able to solve SMD1 and SMD2 but not SMD3 and SMD4, one would infer that the algorithm is unable to handle lower level multi-modality. Such information will be useful for an algorithm developer, as it helps him to identify the specific weaknesses in his approach, which he needs to improve on.

5 Baseline Solution Methodology

In this section, we describe the solution methodology used to solve the constructed test problems. The suggested procedure is a nested bilevel evolutionary algorithm, and requires that a lower level optimization task be solved for every new set of upper level variables produced using the genetic operators. The method relies on a steady state single objective real coded genetic algorithm to solve the problems at both levels.

Table 3: Properties of SMD test problems.

SMD	Upper Level				Lower Level					Conflict
	Constrained	Scalability		Multi-modality	Constrained	Scalability		Multi-modality	Multiple Global Solutions	
		Variables	Constraints			Variables	Constraints			
1	N	Y	-	N	N	Y	-	N	N	N
2	N	Y	-	N	N	Y	-	N	N	Y
3	N	Y	-	N	N	Y	-	Y	N	N
4	N	Y	-	N	N	Y	-	Y	N	Y
5	N	Y	-	N	N	Y	-	Y	N	Y
6	N	Y	-	N	N	Y	-	N	Y	Y
7	N	Y	-	Y	N	Y	-	N	N	Y
8	N	Y	-	Y	N	Y	-	Y	N	Y
9	Y	Y	N	N	Y	Y	N	N	N	Y
10	Y	Y	Y	N	Y	Y	Y	N	N	Y
11	Y	Y	Y	N	Y	Y	N	N	Y	Y
12	Y	Y	Y	N	Y	Y	Y	N	Y	Y

We have implemented a modified version of the procedures [16, 15], which is used to handle the bilevel test problems. The proposed method is based on a single objective Parent Centric Crossover (PCX) [8]. A step-by-step procedure for the algorithm is described as follows:

5.1 Upper Level Optimization Procedure

Step 1: Initialization Scheme. Initialize a random population (N_p) of upper level variables. For each upper level population member execute a lower level optimization procedure to determine the corresponding optimal lower level variables. Assign upper level fitness based on the upper level function value and constraints.

Step 2: Selection of upper level parents. Choose 2μ population members from the previous population and conduct a tournament selection to determine μ parents.

Step 3: Evolution at the upper level. Perform a PCX based crossover [8] (Refer Sub-section 5.4) and a polynomial mutation to create λ offsprings. This provides the upper level variables for each offspring.

Step 4: Lower level optimization. Solve the lower level optimization problem (Refer Sub-section 5.2) for each offspring. This provides the lower level variables for each offspring.

Step 5: Evaluate offsprings. Combine the upper level variables with the corresponding optimal lower level variables for each offspring. Evaluate all the offsprings based on upper level function value and constraints.

Step 6: Population update. Choose r random members from the parent population and pool them with the λ offsprings. The best r members from the pool replace the chosen r members from the population.

Step 7: Termination check. Proceed to the next generation (Step 2) if the termination check (Refer Sub-section 5.6) is false.

5.2 Lower Level Optimization Procedure

The lower level optimization procedure is similar to the upper level procedure except the initialization step which differs slightly. In the following, we provide the steps involved during the lower level optimization task. Let the lower level population size be n_p and the upper level member being optimized be \mathbf{x}_u^0 .

Step 1: If the execution is transferred from Step 1 of the upper level optimization task then go to (a) otherwise go to (b),

a: Initialize n_p lower level member randomly, and assign lower level fitness based on the lower level function value and constraints. Go to Step 2.

b: Initialize $n_p - 1$ lower level members randomly. Determine the member closest to \mathbf{x}_u^0 in the upper level population. The lower level optimal variables from the closest upper level member becomes the n_p^{th} member in the lower level population. Assign lower level fitness based on the lower level function value and constraints. Go to Step 2.

Step 2: Choose 2μ members randomly from the lower level population. Perform a tournament selection with respect to lower level fitness to generate μ parents.

Step 3: Perform crossover and mutation to generate λ offsprings.

Step 4: Evaluate each offspring with respect to lower level function and constraints.

Step 5: Choose r members randomly from the lower level population and pool them with the λ lower level offsprings. The best r members with respect to lower level fitness replace the chosen r members from the lower level population.

Step 6: Proceed to the next generation (Step 2) if the termination check (Refer Sub-section 5.6) is false.

5.3 Parameters

The parameters in the algorithm were fixed as $\mu = 3$, $\lambda = 3$ and $r = 2$. Probability of crossover was fixed as 0.9 and the probability of mutation was fixed as 0.1.

5.4 Crossover Operator

The crossover operator used at both levels is similar to the PCX operator proposed in [15] with minor modifications. The operator creates an offspring from three parents, when one of the three parents is chosen as the index parent.

$$\mathbf{c} = \mathbf{x}_p + \omega_\xi \mathbf{d} + \omega_\eta \frac{\mathbf{p}_2 - \mathbf{p}_1}{2} \quad (52)$$

The terms used in the above equation are defined as follows:

- \mathbf{x}_p is the *index* parent
- $\mathbf{d} = \mathbf{x}_p - \mathbf{w}$, where \mathbf{w} is the mean of μ parents
- \mathbf{p}_1 and \mathbf{p}_2 are the other two parents
- $\omega_\xi = 0.1$ and $\omega_\eta = \sum_{i=1}^{m_v} \frac{m_v}{|x_p^i - w^i|}$ are the two parameters, where $v \in \{u, l\}$ such that m_u is the number of variables at the upper level and m_l is the number of variables at the lower level.

The two parameters ω_ξ and ω_η , describe the extent of variations along the respective directions. While creating $\lambda = 3$ offsprings from $\mu = 3$ parents, each parent is chosen as an index parent at a time.

5.5 Constraint Handling

We define the the constraint violation as the sum of violations of all the constraints at the respective levels. If a member at a particular level has a smaller constraint violation, then it is always preferred over a member with a higher constraint violation at the same level. A member with no constraint violation is deemed to be a feasible, and is considered better than any of the other infeasible members. While comparing two feasible members, the member with a smaller function value at the level is preferred.

5.6 Termination Check

The algorithm uses a variance based termination criteria at both levels. When the value of α_j , described in the following equation becomes less than α_{stop} , the optimization task terminates. In the following, we state the termination criteria at the lower level, which can be similarly extended to the upper level. Let the variance of the lower level population members at generation j for each lower level variable i be v_j^i . If the number of lower level variables is m_l , then α is computed as:

$$\alpha_j = \sum_{i=1}^{m_l} \frac{v_j^i}{v_0^i}. \quad (53)$$

The value of α_j usually lies between 0 and 1 in (53). In the above equation, v_0^i denotes the variance for the variable i in the initial lower level population. For the lower level, the value of α_{stop} is set as 10^{-5} , and for the upper level the value of α_{stop} is set as 10^{-4} .

6 Results

In this section, we provide the results obtained from solving the proposed test problems using the bilevel evolutionary algorithm. The described nested bilevel evolutionary algorithm is a naive scheme, and any intelligent bilevel approach should be expected to produce better results with lesser computational expense. The results are intended as benchmark, and the performance of other schemes may be compared in terms of percentage saving obtained when compared to the proposed nested scheme. We performed 11 number of runs for each of the test problems with 5 and 10 dimensions. In case of 5 dimensions, for SMD1 to SMD5 and SMD7 to SMD12 we choose $p = 1$, $q = 2$ and $r = 1$, and for SMD6 we choose $p = 1$, $q = 0$, $r = 1$ and $s = 2$. In case of 10 dimensions, for SMD1 to SMD5 and SMD7 to SMD12 we choose $p = 3$, $q = 3$ and $r = 2$, and for SMD6 we choose $p = 3$, $q = 1$, $r = 2$ and $s = 2$. The upper level population size N_p and the lower level population size n_p were chosen as 30 for the five dimensional case. Both population sizes were chosen as 50 for the 10 dimensional case.

Results for five dimensional test problems are reported in Tables 4 and 5. Table 4 provides the best, median, and worst values of function evaluations at upper and lower levels. The accuracy achieved and the number of times lower level optimization was performed in a single execution of the bilevel optimization run are reported in Table 5. Similar results for ten dimensional test problems are reported in Tables 6 and 7.

Table 4: Function evaluations (FE) for the upper level (UL) and the lower level (LL) from 11 runs for 5 dimensional test problems.

Pr. No.	Best		Median		Worst	
	Total LL FE	Total UL FE	Total LL FE	Total UL FE	Total LL FE	Total UL FE
SMD1	256858	438	375488	668	582770	1008
SMD2	196744	380	332197	628	613221	1102
SMD3	262703	488	315598	604	439316	844
SMD4	259486	420	366294	608	480675	796
SMD5	222078	444	457265	930	610108	1232
SMD6	334763	540	427114	696	585358	936
SMD7	246375	468	333629	652	685029	1342
SMD8	443430	812	582583	1008	1218196	2076
SMD9	183231	330	284648	514	395735	696
SMD10	179986	480	277696	758	501639	1316
SMD11	11489609	4348	13408524	5086	20540610	7764
SMD12	6211173	354	12950512	738	20983708	1196

The nested bilevel evolutionary algorithm was able to solve all the test problems with five dimensions. We consider a test problem solved if the difference between the function value achieved by the algorithm and the optimal function value is no more than 0.1. However, the number of function evaluations required to obtain the optimal solutions in each of the test problems is large. The function evaluations at the upper level are much smaller, as compared to the function evaluations at the lower level. A large number of lower level function evaluations are required, as a lower level optimization task is executed for each upper level vector. For every newly created upper level vector, we first find the lower level optimal solution and then evaluate the upper level function value. Therefore, the number of function evaluations at the upper level is same as the number of times the lower level optimization task is executed. When the size of the test problems is increased to 10. We observe that the number of function evaluations increase significantly. The nested approach is able to successfully solve the first 5 test problems in all the runs. For test problems SMD6, SMD7 and SMD8, it is unable to solve the problems in all the runs, rather it arrives at the optimal solutions for more than 50% of the runs. For SMD6 the success rate was 87%, for SMD7 it was 66% and for SMD8 it was 62%. The nested approach fails to handle the constrained test problems for the chosen algorithm parameters. The lower level problems could not be completely solved for SMD9 to SMD12, which introduced infeasible members at the upper level.

Table 5: Accuracy for the upper and lower levels, and the lower level calls from 11 runs for 5 dimensional test problems.

Pr. No.	Median	Median	Median	$\frac{\text{LL Evals}}{\text{LL Calls}}$
	UL Accuracy	LL Accuracy	LL Calls	
SMD1	0.000114	0.000087	668	563.89
SMD2	0.000073	0.000016	628	533.01
SMD3	0.000054	0.000055	604	536.47
SMD4	0.000023	0.000057	608	607.80
SMD5	0.000002	0.000009	930	507.82
SMD6	0.000108	0.000061	696	604.64
SMD7	0.000016	0.000177	652	533.84
SMD8	0.000174	0.000027	1008	562.69
SMD9	0.000017	0.000054	514	553.54
SMD10	0.034759	0.018510	758	367.04
SMD11	0.0131643	0.0129893	5086	2635.64
SMD12	0.032372	0.000206	738	19202.32

The results demonstrate that a high number of function evaluations are required to solve bilevel problems. With an increase in the number of dimensions, the complexity increases significantly and the available computational resources quickly become insufficient to solve larger versions of the problems. In this paper, we utilize a global optimizer at both levels, which successfully solved smaller versions of the test problems, but failed for constrained test problems with high dimensions. Given, the complex nature of bilevel optimization problems, evolutionary algorithms might be a useful approach to follow. However, using evolutionary algorithms alone would demand a large number of function evaluations to solve even simple bilevel problems. Therefore, an intelligent approach which utilizes results from the classical literature within an evolutionary algorithm might be a feasible direction towards handling such problems. The set of test problems proposed in this paper would be useful to evaluate such algorithms across various difficulties which a bilevel optimization problem could offer.

7 Conclusions

In this paper we have provided a test problem construction procedure for unconstrained as well as constrained bilevel optimization. The construction procedure offers the flexibility to control the difficulties at the two levels individually as well as collectively. The construction procedure has been used to create a test-bed of 12 bilevel optimization problems, out of which 8 are unconstrained and 4 are constrained. The test-suite contains problems, which are scalable in terms of number of variables as well as constraints. Moreover, the optimal solutions for all the test problems are clearly identified, which would be useful in testing and evaluating bilevel optimization algorithms.

Table 6: Function evaluations (FE) for the upper level (UL) and the lower level (LL) from 11 runs for 10 dimensional test problems. A ‘x’ denotes that the algorithm terminated far away ($\Delta F \geq 0.1$) from the optimal solution. A ‘-’ denotes that a feasible solution could not be obtained for the test problem.

Pr. No.	Best		Median		Worst	
	Total LL FE	Total UL FE	Total LL FE	Total UL FE	Total LL FE	Total UL FE
SMD1	862653	1080	1623356	2534	2074334	3488
SMD2	1055976	1398	1467246	2366	2114442	3418
SMD3	900358	1210	1383632	2278	1805562	2862
SMD4	566344	678	1087632	1598	1314986	2028
SMD5	1226344	1620	1993124	2890	2483442	3492
SMD6	1225742	1502	2224450	2936	3786498 (x)	4278 (x)
SMD7	932460	1382	1566481	2394	2435994 (x)	3858(x)
SMD8	1457480	2116	2710132	4188	5294734 (x)	5986 (x)
SMD9	-	-	-	-	-	-
SMD10	-	-	-	-	-	-
SMD11	-	-	-	-	-	-
SMD12	-	-	-	-	-	-

The test problem construction procedure should allow researchers to create additional test problems by varying the basic functions used in different test problems. As a benchmark for comparison, we have provided results from a nested bilevel evolutionary scheme, which utilizes a global optimizer at both levels. Five and ten-variable instances of all the test problems have been solved, which demonstrate the high computational requirement of bilevel problems even for smaller instances. This imply indicates that the solution of bilevel problems, even with an evolutionary algorithm, is a challenging task and more attention must now be made to develop computationally faster algorithms.

Acknowledgments

Authors A. Sinha and P. Malo wish to thank the Wallenberg foundation and Liikesivistysrahasto for supporting this study. K. Deb and A. Sinha acknowledge the support provided by the Academy of Finland under grant number 133387.

References

- [1] E. Aiyoshi and K. Shimizu. Hierarchical decentralized systems and its new solution by a barrier method. *IEEE Transactions on Systems, Man, and Cybernetics*, 11:444–449, 1981.

Table 7: Accuracy for the upper and lower levels, and the lower level calls from 11 runs for 10 dimensional test problems. A - denotes that a feasible solution could not be obtained for the test problem.

Pr. No.	Median	Median	Median	$\frac{\text{LL Evals}}{\text{LL Calls}}$
	UL Accuracy	LL Accuracy	LL Calls	
SMD1	0.000332	0.000018	2534	644.54
SMD2	0.000066	0.000011	2366	653.36
SMD3	0.000359	0.000033	2278	655.76
SMD4	0.000286	0.000027	1598	685.43
SMD5	0.000052	0.000009	2890	716.82
SMD6	0.001435	0.000082	2936	768.34
SMD7	0.006263	0.000127	2394	654.34
SMD8	0.003122	0.000157	4188	647.12
SMD9	-	-	-	-
SMD10	-	-	-	-
SMD11	-	-	-	-
SMD12	-	-	-	-

- [2] J.F. Bard. *Practical Bilevel Optimization: Algorithms and Applications*. The Netherlands: Kluwer, 1998.
- [3] J.F. Bard and J. Falk. An explicit solution to the multi-level programming problem. *Computers and Operations Research*, 9:77–100, 1982.
- [4] L. Bianco, M. Caramia, and S. Giordani. A bilevel flow model for hazmat transportation network design. *Transportation Research Part C: Emerging technologies*, 17(2):175–196, 2009.
- [5] P. H. Calamai and L. N. Vicente. Generating linear and linear-quadratic bilevel programming problems. *SIAM J. Sci. Comput.*, 14(1):770–782, 1992.
- [6] P. H. Calamai and L. N. Vicente. Generating quadratic bilevel programming test problems. *ACM Trans. Math. Software*, 20(1):103–119, 1994.
- [7] B. Colson, P. Marcotte, and G. Savard. An overview of bilevel optimization. *Annals of Operational Research*, 153:235–256, 2007.
- [8] K. Deb, A. Anand, and D. Joshi. A computationally efficient evolutionary algorithm for real-parameter optimization. *Evolutionary Computation Journal*, 10(4):371–395, 2002.
- [9] K. Deb and A. Sinha. An efficient and accurate solution methodology for bilevel multi-objective programming problems using a hybrid evolutionary-local-search algorithm. *Evolutionary Computation Journal*, 18(3):403–449, 2010.

- [10] S. Dempe. Annotated bibliography on bilevel programming and mathematical programs with equilibrium constraints. *Optimization*, 52(3):339–359, 2003.
- [11] S. Dempe, J. Dutta, and S. Lohse. Optimality conditions for bilevel programming problems. *Optimization*, 55:505–524, 2006.
- [12] J. Herskovits, A. Leontiev, G. Dias, and G. Santos. Contact shape optimization: A bilevel programming approach. *Struct Multidisc Optimization*, 20:214–221, 2000.
- [13] A. Mitsos and P. I. Barton. A test set for bilevel programs, 2006. <http://yoric.mit.edu/download/Reports/bileveltestset.pdf>.
- [14] K. Moshirvaziri, M. A. Amouzegar, and S. E. Jacobsen. Test problem construction for linear bilevel programming problems. *Journal of Global Optimization*, 8(3):235–243, 1996.
- [15] A. Sinha, A. Srinivasan, and K. Deb. A population-based, parent centric procedure for constrained real-parameter optimization. In *2006 IEEE Congress on Evolutionary Computation (CEC-2006)*, pages 239–245. IEEE Press, 2006.
- [16] A. Sinha, S. Tiwari, and K. Deb. A population-based, steady-state procedure for real-parameter optimization. In *2005 IEEE Congress on Evolutionary Computation (CEC-2005)*, pages 514–521. IEEE Press, 2005.
- [17] L. N. Vicente and P. H. Calamai. Bilevel and multilevel programming: A bibliography review. *Journal of Global Optimization*, 5(3):291–306, 2004.
- [18] G. Wang, Z. Wan, X. Wang, and Y. Lv. Genetic algorithm based on simplex method for solving linear-quadratic bilevel programming problem. *Comput Math Appl*, 56(10):2550–2555, 2008.
- [19] Y. Yin. Genetic algorithm based approach for bilevel programming models. *Journal of Transportation Engineering*, 126(2):115–120, 2000.